arXiv:1204.2157v1 [quant-ph] 10 Apr 2012

# Decoherence Dynamics of Measurement-Induced Nonlocality and comparison with Geometric Discord

Ajoy Sen,[1,*] Debasis Sarkar,[1,†] and Amit Bhar[2,‡]
[1]*Department of Applied Mathematics, University of Calcutta, 92, A.P.C. Road, Kolkata-700009, India*
[2]*Department of Mathematics, Jogesh Chandra Chaudhuri College,*
*30, Prince Anwar Shah Road, Kolkata-700033, India*

(Dated: March 2, 2013)

We check the decoherence dynamics of Measurement-induced Nonlocality and compare it with geometric discord. There are quantum states, on which the action of dephasing channel cannot destroy MIN in finite or infinite time. We also investigate the additive dynamics of MIN on a qubit state under two independent noise. Geometric discord also follows such additive dynamics like quantum discord.

PACS numbers: 03.67.Mn, 03.65.Yz.

## I. INTRODUCTION

Separability, Nonlocality and quantumness these are the three most discussed areas in Quantum Information theory. While the first one has a definite identifier in terms of entanglement measures; the last two lacks definite identifiers as they can exist in various forms. But there is one thing common among all of them. They all help to understand the state space structures of quantum systems from different perspectives. The idea behind discord was to differentiate classicality from quantumness[1, 2]. There are separable states which have non-zero value of it, indicating existence of quantumness in separable states. Measurement-induced nonlocality(MIN) as introduced by Luo and Fu[3] is another type of correlation which reveals a type of nonlocality that exists in state space. There are classical states(i.e., vanishing discord) with non-zero MIN. Since our classical world is local so existence of non-zero MIN for classical states exhibits a kind of non-classical trait in those states.

Study on the behavior of non-classical correlations under decoherence started with the strange observation of Entanglement sudden death(ESD)[4]. For some class of initial states ESD is observed under independent Markovian decoherence but similar behavior is not observed in case of discord [5]. In fact quantum discord decays monotonically under such decoherences and sudden death of discord can not occur under Markovian decoherence[6]. Yu and Eberly had also pointed out non-additivity of decoherence channel under entanglement dynamics but opposite result is observed in case of quantum discord. The dynamics of quantum discord under decoherence channels and Pauli channels is also well studied in [5, 7]. Geometric discord(GD), as introduced by Dakic and Vedral[2] is a way to visualize quantum discord geometrically. Behavior of different measures of discord can be quite different[8]. The value of Geometric discord matches with the value of MIN for pure 2 qubit states. Here we will investigate the decoherence dynamics of MIN under dissipative(depolarizing, amplitude damping) and non-dissipative(dephasing) channels. We will take three kinds of initial states and compare the dynamics with the corresponding dynamics of geometric discord. We will also investigate the additivity of decoherence channels under MIN and GD dynamics. In our study, we will use original version of MIN as introduced by Luo and Fu. However there are entropic formulation of MIN. Few results regarding this area can be found in [9–12].

## II. PRELIMINARY NOTIONS ON MIN AND GEOMETRIC DISCORD

Let $\rho$ be any bipartite state shared between two parties A and B. Then MIN (denoted by $N(\rho)$) is defined as[3],

$$N(\rho) := \max_{\Pi^A} \parallel \rho - \Pi^A(\rho) \parallel^2 \tag{1}$$

where maximum is over all von Neumann measurements $\Pi^A$ which do not disturb $\rho_A$, the local density matrix of A, i.e., $\Sigma_k \Pi_k^A \rho_A \Pi_k^A = \rho_A$ and $\|.\|$ is taken as Hilbert Schmidt norm (i.e. $\parallel X \parallel = [Tr(X^\dagger X)]^{\frac{1}{2}}$). On the other hand Geometric discord for a quantum state $\rho$ is defined as[2],

$$D_g(\rho) = \min_{\chi \in \Omega_0} \parallel \rho - \chi \parallel^2 \tag{2}$$

where $\Omega_0$ is the set of all zero discord states and $\parallel . \parallel$ is the usual Hilbert-Schmidt norm. Any state $\rho_{AB}$ of a $2 \otimes 2$ dimensional quantum system can be expressed in

---

[*] Electronic address: ajoy.sn@gmail.com
[†] Electronic address: dsappmath@yahoo.com
[‡] Electronic address: bhar.amit@yahoo.com

the form

$$\rho_{AB} = \frac{1}{\sqrt{4}}\frac{I^A}{\sqrt{2}} \otimes \frac{I^B}{\sqrt{2}} + \sum_{i=1}^{3} x_i X_i^A \otimes \frac{I^B}{\sqrt{2}} + \frac{I^A}{\sqrt{2}} \otimes \sum_{i=1}^{3} y_i Y_i^B + \sum_{i=1}^{3}\sum_{j=1}^{3} t_{ij} X_i^A Y_j^B \tag{3}$$

where $\{X_i^A : \text{i=0,1,2,3}\}$ and $\{Y_j^B : j = 0, 1, 2, 3\}$ are the orthonormal Hermitian operator bases for $L(H^A)$ and $L(H^B)$ respectively with $X_0^A = Y_0^B = I^A/\sqrt{2}$. Here we will use the Pauli matrices (with normalization factor $\frac{1}{\sqrt{2}}$) as the operator base. For this state(3) MIN[3] and geometric discord[2] can be written as

$$N(\rho_{AB}) = \begin{cases} trTT^t - \frac{1}{\|\mathbf{x}\|^2}\mathbf{x}^t TT^t \mathbf{x} & \text{if } \mathbf{x} \neq \mathbf{0} \\ trTT^t - \lambda_{min} & \text{if } \mathbf{x} = \mathbf{0} \end{cases} \tag{4}$$

$$D_g(\rho_{AB}) = (\| \overrightarrow{x} \|^2 + \| T \|^2 - \lambda_{max}) \tag{5}$$

where the matrix $T = (t_{ij})_{3\times3}$ is the correlation matrix with $\lambda_{min}$ being minimum eigenvalue of $TT^t$ and $\|\mathbf{x}\|^2 := \sum_i x_i^2$ with Bloch vector $\mathbf{x} = (x_1, x_2, x_3)^t$. $\lambda_{max}$ is the maximum eigenvalue of $xx^t + TT^t$. The elements of the correlation matrix and Bloch vector can be obtained from the following relations,

$$\begin{aligned} t_{ij} &= tr(\rho\frac{\sigma_i}{\sqrt{2}} \otimes \frac{\sigma_j}{\sqrt{2}}), \quad i, j = 1, 2, 3 \\ x_i &= tr(\rho\frac{\sigma_i}{\sqrt{2}} \otimes \frac{I}{\sqrt{2}}), \quad i = 1, 2, 3 \end{aligned} \tag{6}$$

## III. EVOLUTION OF MIN AND GEOMETRIC DISCORD

Let us consider two qubits, interacting independently with the individual environments. Then their evolution can be described by Lindblad equation. This can be written in terms of Kraus operators in the form

$$\rho(t) = \sum_{\mu,\nu} E_{\mu,\nu}\rho(0)E_{\mu,\nu}^{\dagger} \tag{7}$$

where $\rho(0)$ is the initial state, $\rho(t)$ denotes the state after time $t$ and $E_{\mu,\nu}$ are the Kraus operators which satisfies the relation $\sum_{\mu,\nu} E_{\mu,\nu}^{\dagger}E_{\mu,\nu} = I$ for all $t$.
For studying the evolution of correlations, here we will consider three kinds of initial states

$$|\psi\rangle = \sqrt{1-\alpha}|00\rangle + \sqrt{\alpha}|11\rangle, \alpha \in [0, 1] \tag{8}$$

$$\rho_2 = \frac{1-\alpha}{4}I + \alpha|\psi^-\rangle\langle\psi^-|, \alpha \in [0, 1] \tag{9}$$

$$\rho_3 = \frac{\alpha}{2}(|00\rangle\langle00| + |11\rangle\langle11|) + (1-\alpha)|\psi^-\rangle\langle\psi^-|, \alpha \in [0, 1] \tag{10}$$

The first one is a pure state, second one is a Werner state and third one is a mixture of a mixed state with the singlet state($|\psi^-\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$). These three kinds of states are basically X states. The X state structure of theese three states remains unchanged under the action of the decoherence channels. We will consider that the both qubit decays at same rate under each decoherence channel.

**Depolarizing Channel**: Application of this type of channel depolarize the density matrix to completely mixed state with probability $p$. In this case the Kraus operators are of the form: $E_0 = \sqrt{1-\frac{3\gamma}{4}}I, E_1 = \sqrt{\frac{\gamma}{2}}X, E_2 = \sqrt{\frac{\gamma}{2}}Y, E_3 = \sqrt{\frac{\gamma}{2}}Z$ and $(X, Y, Z) \equiv (\sigma_x, i\sigma_y, \sigma_z)$. Consider the Initial state $\rho(0) = |\psi\rangle\langle\psi|$ where $|\psi\rangle$ is given in (8). Under the independent action of the channel on both qubit the density matrix elements evolve as,

$$\begin{aligned} \rho_{11}(t) &= \rho_{11}(0)(1-\gamma) + \frac{\gamma^2}{4} \\ \rho_{22}(t) &= \rho_{33}(t) = \frac{\gamma}{2}(1-\frac{\gamma}{2}) \\ \rho_{44}(t) &= 1 - \rho_{11}(t) - 2\rho_{22}(t) \\ \rho_{14}(t) &= \rho_{41}(t) = \rho_{14}(0)(1-\gamma)^2 \end{aligned} \tag{11}$$

the evolution of Bloch vector and correlation matrix can be obtained from the relations in (6). MIN and Geometric discord for this initial state is obtained as

$$N(\rho(t)) = D_g(\rho(t)) = 2\alpha(1-\alpha)(1-\gamma)^4 \tag{12}$$

If the initial condition be of the form (9) then noting that $\|x(t)\| = 0, \forall t$ we have

$$D_g(\rho(t)) = N(\rho(t)) = \frac{\alpha^2}{2}(1-\gamma)^4 \tag{13}$$

If we consider $\gamma = 1 - Exp[-\Gamma_1 t]$ i.e., $\Gamma_1$ is the rate of depolarizing then we observe monotonic nature of decay of MIN and geometric discord as $\gamma \to 1 (\equiv t \to \infty)$.

**Dephasing Channel**: Physically, dephasing corresponds to any process of losing coherence without any exchange of energy. Kraus operators corresponding to this process are given by

$$E_0 = \begin{bmatrix} 1 & 0 \\ 0 & \sqrt{1-\gamma} \end{bmatrix}, E_1 = \begin{bmatrix} 0 & 0 \\ 0 & \sqrt{\gamma} \end{bmatrix}$$

Under the action of this channel the density matrix elements corresponding to the initial state (8) evolve as

$$\begin{aligned} \rho_{ii}(t) &= \rho_{ii}(0) \quad \text{for all} \quad i = 1, 2, 3, 4 \\ \rho_{14}(t) &= \rho_{41}(t) = \rho_{14}(0)(1-\gamma) \end{aligned} \tag{14}$$



FIG. 1: *Monotonic nature of decay of MIN for the pure qubit initial state under independent depolarising noise. For* $\alpha = 0$ *or* 1 *MIN remains zero always.*

MIN and discord for this state is given by

$$N(\rho(t)) = \begin{cases} \frac{1}{4} + \frac{(1-\gamma)^2}{4} & \text{if } \alpha = \frac{1}{2} \\ 2\alpha(1-\alpha)(1-\gamma)^2 & \text{if } \alpha \neq \frac{1}{2} \end{cases} \quad (15)$$

$$D_g(\rho(t)) = 2\alpha(1-\alpha)(1-\gamma)^2$$

we take $\gamma = 1 - Exp[-\Gamma_2 t]$ where $\Gamma_2$ is the rate of dephasing.



FIG. 2: *The nature of decay of MIN and geometric discord and respectively for the pure initial state under independent dephasing noise. While the value of discord decays monotonically for all* $\alpha$*, MIN for* $\alpha = 0.5$ *does not vanishes even in infinite time.*

Again if the initial state be of the form (9) then

$$\begin{aligned} \rho_{ii}(t) &= \rho_{ii}(0) \quad \text{for all} \quad i = 1,2,3,4 \\ \rho_{23}(t) &= \rho_{32}(t) = \rho_{23}(0)(1-\gamma) \end{aligned} \quad (16)$$

MIN and discord for this state is given by

$$\begin{aligned} N(\rho(t)) &= (\frac{\alpha}{2})^2(1-\gamma)^2 + (\frac{\alpha}{2})^2 \\ D_g(\rho(t)) &= 2(\frac{\alpha}{2})^2(1-\gamma)^2 \end{aligned} \quad (17)$$

From Fig.2 we see that MIN for maximally entangled state does not vanishes as $\gamma \to 1$. So under dephasing noise while the quantumness vanishes, nonlocality remains present throughout the process for this state. In this sense nonlocality is more robust than quantumness. Similar kind of conclusion can be obtained for the Werner state(Fig.3). In this case for any non-zero $\alpha$ MIN does not decay to zero like geometric discord. Decay of MIN in case of Werner states clearly suggests that Werner state is better candidate than pure state for preserving MIN in any dephasing process.



FIG. 3: *The nature of decay of MIN and geometric discord and respectively for the Werner state under independent dephasing noise. MIN shows more robustness against dephasing noise than geometric discord.*

**Generalized Amplitude Damping Channel**: Exchange of energy occurs in this type of decoherence noise. Kraus operators corresponding to this channel are given by

$$E_0 = \sqrt{p}\begin{bmatrix} 1 & 0 \\ 0 & \sqrt{1-\gamma} \end{bmatrix}, E_1 = \sqrt{p}\begin{bmatrix} 0 & \sqrt{\gamma} \\ 0 & 0 \end{bmatrix}$$

$$E_2 = \sqrt{1-p}\begin{bmatrix} \sqrt{1-\gamma} & 0 \\ 0 & 0 \end{bmatrix}, E_3 = \sqrt{1-p}\begin{bmatrix} 0 & 0 \\ \sqrt{\gamma} & 0 \end{bmatrix}$$

Evolution of the matrix element for the initial state (8) can be written as

$$\begin{aligned} \rho_{11}(t) =& (1-(1-p)\gamma)^2 - \alpha(1-\gamma)(1-(1-2p)\gamma) \\ \rho_{22}(t) =& -\gamma(-1+\alpha+p(1-2\alpha(1-\gamma)-2\gamma)+\gamma \\ & (1+p^2-\alpha)) \\ \rho_{33}(t) =& \rho_{22}(t) \\ \rho_{44}(t) =& 1-\rho_{11}(t)-2\rho_{22}(t) \\ \rho_{14}(t) =& \sqrt{\alpha-\alpha^2}(1-\gamma) \end{aligned} \quad (18)$$

and the dynamics of the density matrix elements for the initial state (9) is given by

$$\begin{aligned}\rho_{11}(t) =&\frac{1}{4}(-\alpha(1-\gamma)^2+(1+(2p-1)\gamma)^2)\\ \rho_{22}(t) =&\rho_{33}(t)=\frac{1}{4}(1+\alpha(1-\gamma)^2-(1-2p)^2\gamma)^2)\\ \rho_{44}(t) =&1-\rho_{11}(t)-2\rho_{22}(t)\\ \rho_{23}(t) =&\frac{\alpha}{2}(1-\gamma)\end{aligned} \tag{19}$$

We plot the figures corresponding to the initial state (8) and (9) taking three values of $p$ and $\gamma = 1 - Exp[-\Gamma_3 t]$. $\Gamma_3$ is the rate of damping. $p = 1$ corresponds to usual amplitude damping channel. We also compare the dynamics with the corresponding dynamics of geometric discord in the figures(Fig.4, Fig.5). From the figures, we observe monotonic nature of decay of MIN like geometric discord. No sudden death type behavior for MIN occurs. Clearly MIN does not disappear in finite time. While the decay in the case of MIN is smooth, the corresponding decay in case of discord is not so smooth. There are certain regions where we can notice changing behavior of discord. Another important observation is that the application of amplitude damping completely decays the correlations as $\gamma \to 1$ for both the initial states. In case of pure state when $\alpha = 0$ or 1 MIN remains zero always.



FIG. 4: *Monotonic nature of decay of geometric discord and MIN respectively for the Werner state under independent generalized amplitude damping for* $p = 1, 0.5, 0.67$ *respectively. The white curve in the figures of discord shows its changing nature while the decay of MIN remains smooth.*



FIG. 5: *Monotonic nature of decay of geometric discord and MIN respectively for the pure state under independent generalized amplitude damping for* $p = 1, 0.5, 0.67$ *respectively. Discord shows its changing nature while MIN remains smooth.*

**Dephasing and Amplitude damping**: Now let us consider the scenario where both qubits are acted simultaneously by both dephasing and amplitude damping noise. Since the diagonal elements remains unchanged under dephasing noise, so, in this case the off diagonal elements decay as the sum of the individual rate of decay. Here we consider that the both channel decays at the same rate $\Gamma_2 = \Gamma_3 = \Gamma$. Under this consideration the dynamics of density matrix elements for the initial state (8) is

$$\begin{aligned}\rho_{11}(t) =&1-\alpha+\alpha(1-\exp[-\Gamma t])^2\\ \rho_{22}(t) =&\alpha(1-\exp[-\Gamma t])^2\exp[-\Gamma t]\\ \rho_{33}(t) =&\rho_{22}(t)\\ \rho_{44}(t) =&\alpha\exp[-2\Gamma t]\\ \rho_{23}(t) =&\sqrt{(\alpha-\alpha^2)}\exp[-2\Gamma t]\end{aligned} \tag{20}$$

In Fig.6, we plot the values of MIN and compared with discord. The nature of decay reveals additivity in the nature of decay of MIN under simultaneous action of both the noise. Discord also shows the same behavior. We also compare the rate of decay of MIN for three types of decoherence channels, described earlier, for the Bell state. From Fig.8 we observe that MIN is more robust under dephasing noise than other two types of noise. In fact dephasing noise can not fully destroy this correlation in infinite run.

So far we have discussed dissipative dynamics for initial states $|\psi\rangle$ (8) and $\rho_2$(9). Similar type of analysis can also be done by taking the initial state $\rho_3$(10). In the Fig.7 we plot the decay nature corresponding to this initial state $\rho_3$ for a particular $\alpha$. From the figure we observe that the dephasing noise can not decay MIN



FIG. 6: *Monotonic nature of decay geometric discord and MIN respectively for pure qubit state assuming simultaneous action of both dephasing and amplitude damping channels($p = 1$) in each qubit.*



FIG. 7: *Comparison of the rate of decay of MIN and discord under three types of decoherence channels for initial state $\rho_3$ at $\alpha = 0.25$*

to zero in this case also. But there is a particular time when discord and MIN matches exactly and after that discord falls faster and ultimately goes to zero. Under dephasing noise discord always remains below than MIN and they both decays to zero. In case of amplitude damping noise discord falls slowly than MIN and at one time their values become same and ultimately they both decay at the same rate.



FIG. 8: *Comparison of the rate of decay of MIN for the Bell state $|\phi^+\rangle$ under the three decoherence channels.*

## IV. CONCLUSION

Thus we have compared two different types of correlations of quantum system under decoherence dynamics. We have found additive dynamics of MIN under simultaneous action of decoherence noise. Geometric discord also behaves similarly. We have also found some states which retain their nonlocality even after applying a type of decoherence noise. Such behavior is not observed for geometric discord for those states. Under amplitude damping noise sudden change in the decay of geometric discord is also observed but in this case MIN decays smoothly without any change. After completing our work we recently observed another work in this line [13].

**Acknowledgement.** The author A. Sen acknowledges the financial support from University Grants Commission, New Delhi, India.

---

[1] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001);D. Cavalcanti, L. Aolita, S. Boixo, K. Modi, M. Piani,and A. Winter, Phys. Rev. Lett., 106, 220403(2011).
[2] B. Dakic, V. Vedral and C. Brukner, Phys. Rev. Lett. **105**, 190502 (2010).
[3] S. Luo and S. S. Fu, Phys. Rev. Lett. **106**, 120401 (2011).
[4] T. Yu and J.H. Eberly, Phys. Rev. Lett. **97**, 140403 (2006).
[5] T. Werlang et.al., Phys. Rev. A **80**, 024103 (2000).
[6] A. Ferraro, L. Aolita, D. Cavalcanti, F. M. Cucchietti, A. Acin, Phys. Rev. A **81**,052318 (2009).
[7] J. Maziero, L. C. Celeri, R. M. Serra, V. Vedral, Phys.

Rev. A **80**,044102 (2009).

[8] B. Bellomo, R. L. Franco, G. Compagno, arXiv: 1104.4043.

[9] Guo-Feng Zhang, arXiv: 1201.1949v1.

[10] Ming-Liang Hu and Heng Fan,arXiv:1201.6430v2

[11] J. Maziero, L. C. Céleri, R. M. Serra and V. Vedral, Phys. Rev. A **80**, 044102 (2009); T. Werlang, S. Souza, F. F. Fanchini and C. J. Villas Boas, Phys. Rev. A **80**, 024103 (2009); S. Luo, S. Fu and Nan Li, Phys. Rev. A **82**, 052122 (2010).

[12] S. Luo and N. Li, Phys. Rev. A **84**, 052309 (2011); Patrick J. Coles, arXiv:1110.1664; F. Ciccarello and V. Giovannetti , arXiv:1105.5551.

[13] M. Ramzan, arXiv:1204.1900.